

MAT 1033 Bookmark

Slope of a line	$m = \frac{y_2 - y_1}{x_2 - x_1}$
Slope-intercept form	$y = mx + b$
Point-slope form	$y - y_1 = m(x - x_1)$ or $y = m(x - x_1) + y_1$
Slope of relational lines	$m_1 = m_2$; parallel $m_2 = \frac{-1}{m_1}$; perpendicular
Quadratic formula	Given $ax^2 + bx + c = 0$; $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Vertical line	$x = a$; Undefined slope
Horizontal line	$y = b$; Slope = 0
x-intercept	Let $y = 0$; $(f(x) = 0, 0)$
y-intercept	Let $x = 0$; $(0, f(0))$

Geometry Formulas

Area rectangle	$A = L W$
Perimeter of rectangle	$P = 2L + 2W$
Area of circle	$A = \pi r^2$
Circumference of circle	$C = 2\pi r$
Volume of a cube	$V = s^3$ or LWH

Math Translation Words

+ → Sum, increased by, addition, more than	$x \rightarrow$ Product, multiply, of
- → Difference, subtract, decreased by, less than	$\div \rightarrow$ Divide, quotient
= → Equal, is	

*If $AB = 0$, then $A = 0$ or $B = 0$

*If $x^2 = k$, $k > 0$, then $x = \pm \sqrt{k}$

Exponent Rules

$m^a m^b = m^{a+b}$	$(m^a n^c)^b = m^{ab} n^{bc}$
$\frac{m^a}{m^b} = m^{a-b}$	$m^{-a} = \frac{1}{m^a}$
$m^0 = 1$	$\sqrt[b]{m^a} = m^{\frac{a}{b}}$
n even, $\sqrt[n]{a^n} = a $	n odd, $\sqrt[n]{a^n} = a$
$i = \sqrt{-1}$, $i^2 = -1$	$\sqrt{-m} = i\sqrt{m}$

Factoring Summary

GCF:	$3x^2 + 9x + 15 \rightarrow 3(x^2 + 3x + 5)$	
4 terms-grouping	$3x^3 + 2x^2 - 6x - 4 =$ $(3x^3 + 2x^2) + (-6x - 4) =$ $x^2(3x + 2) - 2(3x + 2)$ $\rightarrow (3x + 2)(x^2 - 2)$	
a = 1	$x^2 + 4x - 12$: find factors of -12, add to 4, $\rightarrow (x - 2)(x + 6)$	
$x^2 - y^2$	$(x - y)(x + y)$	
$x^2 + y^2$	Does not factor/prime	
$ax^2 + bx + c$ $a \neq 1$	$3x^2 + 2x - 8$: factors of 3&8 that give difference of 2	$(3, 1)$ & $(1, 2, 4, 8)$ $4 \cdot 1 - 3 \cdot 2 =$ $4 - 6 = -2$ $3x^2 + 2x - 8 \rightarrow$ $(3x - 4)(x + 2)$
Perfect squares	$p^2 \pm 2pq + q^2$: $4x^2 - 12x + 9 \rightarrow (2x - 3)^2$	

Factoring steps when solving quadratic:

1. Get the equation = 0
2. Factor out any common terms
3. Is it a difference of two squares?
4. Does it have 4 terms (grouping)
5. For a trinomial, use AC or trial/error.
6. Set all factors with a variable = 0 and solve.